



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ALGEBRA.

181. Proposed by J. F. LAWRENCE, Breckenridge, Mo.

$$\text{Show that } \phi(1)\frac{x}{1+x^2} - \phi(3)\frac{x^3}{1+x^6} + \phi(5)\frac{x^5}{1+x^{10}} - \dots \text{ ad inf. } = \frac{x(1-x^2)}{(1+x^2)^2} \phi(n)$$

being the number of integers less than n and prime to it. [From Hall and Knight's *Higher Algebra*, page 358].

182. Proposed by J. F. LAWRENCE, Breckenridge, Mo.

Find the values of $x_1, x_2, x_3, \dots, x_n$ which satisfy the following system of simultaneous equations:

$$\frac{x_1}{a_1 - b_1} + \frac{x_2}{a_1 - b_2} + \dots + \frac{x_n}{a_1 - b_n} = 1.$$

$$\frac{x_1}{a_2 - b_1} - \frac{x_2}{a_2 - b_2} + \dots + \frac{x_n}{a_2 - b_n} = 1.$$

$$\dots \dots \dots \dots \dots$$

$$\frac{x_1}{a_n - b_1} + \frac{x_2}{a_n - b_2} + \dots + \frac{x_n}{a_n - b_n} = 1.$$

GEOMETRY.

203. Proposed by W. J. GREENSTREET, A. M., Editor of *The Mathematical Gazette*, Stroud, Eng.

Show that two parabolaes can always be drawn through the vertices of a triangle to touch its circumcircle at a vertex, and that the axes of these pairs of curves are orthogonal. Show that any triangle may be circumscribed by a conic so that the tangents at each vertex is parallel to the opposite side.

204. Proposed by ELMER SCHUYLER, B. Sc., Professor of German and Mathematics, Boys' High School, Reading, Pa.

Construct a triangle, having given an angle, the length of its bisector, and the sum of the including sides. [Phillips and Fisher].

CALCULUS.

168. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

The tangent of what Cartesian curve makes an x -intercept always m times as long as the corresponding y -intercept?

169. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

Find the value of y from the Eulerian equation

$$y = \int \frac{dx}{(x + \sqrt{3})^{\frac{2}{3}}(x^2 + 1)}.$$